

Monomial Golod Quotients of Exterior Algebras

Emil Sköldberg

metadata, citation and similar papers at core.ac.uk

Communicated by Craig Huneke

Received September 4, 1997

It is proved that for any set M of squarefree monomials in the variables x_1, \dots, x_n , the algebra $A = k[x_1, \dots, x_n]/(M)$ is Golod if and only if the algebra $B = E(x_1, \dots, x_n)/(M)$ is Golod, where E is the exterior algebra. This is proved by showing the equivalence of the extremality of the Poincaré series of A and B .

© 1999 Academic Press

1. INTRODUCTION

Let Δ be a simplicial complex on the vertex set $[n] = \{1, 2, \dots, n\}$. To each such Δ one can associate an ideal I_Δ in the polynomial ring $S = k[x_1, x_2, \dots, x_n]$ by letting I_Δ be generated by the set of monomials

$$\{x_{i_1}x_{i_2} \cdots x_{i_m} \mid \{i_1, i_2, \dots, i_m\} \notin \Delta\}. \quad (1)$$

Conversely, every ideal in S generated by squarefree monomials comes from some simplicial complex in this way. The quotient ring S/I_Δ is then called the Stanley–Reisner ring of Δ , and is denoted by $k[\Delta]$. If we denote the exterior algebra over k on the variables x_1, x_2, \dots, x_n by E , we may form the ideal J_Δ in E by letting it be generated by the set (1). The quotient E/J_Δ is now denoted by $k\{\Delta\}$, and is called the indicator algebra of Δ .

The relations between $k[\Delta]$ and $k\{\Delta\}$, or between more general quotients S/I and E/J , have been investigated from several viewpoints, some of which we will recall:

- Macaulay's theorem, which gives an upper bound for the possible Hilbert series of S/I , has a counterpart in the Kruskal–Katona theorem which gives a similar bound for the Hilbert series of E/J .

• Gotzmann's theorem, which essentially says that an equality in Macaulay's bound in a degree will persist in higher degrees, if no new relations are introduced, has a counterpart for quotients of the exterior algebra as well. This is proved by Aramova *et al.* [AHH97].

• The correspondence between S -resolutions of $k[\Delta]$ and E -resolutions of $k\{\Delta\}$ has been studied by among others Aramova *et al.* [AAH]. In particular they give the relation between $\dim_k \operatorname{Tor}_*^S(k[\Delta], k)$ and $\dim_k \operatorname{Tor}_*^E(k\{\Delta\}, k)$.

The purpose of the present paper is to relate the properties of $\operatorname{Ext}_{k[\Delta]}^*(k, k)$ and $\operatorname{Ext}_{k\{\Delta\}}^*(k, k)$ to each other in one aspect, namely to prove that $k[\Delta]$ is Golod if, and only if, $k\{\Delta\}$ is Golod. Since there are classes of Stanley–Reisner rings which are known to be Golod, (e.g., when I_Δ is a squarefree lexsegment ideal [AHH98], or more generally, when I_Δ is squarefree 0-Borel fixed [Pee96], this gives new examples of Golod quotients of exterior algebras.

The author expresses his gratitude to Jürgen Herzog for stimulating discussions on this subject, to Jörgen Backelin for pointing out the existence of Lemma 4, and to the referee for suggesting substantial improvements to the paper, for example by simplifying the proofs of Lemma 4 and Proposition 1.

2. NOTATION AND DEFINITIONS

We will let k be an arbitrary field, and we will consider finitely generated $\mathbb{N} \times \mathbb{Z}_2$ -graded k -algebras. All algebras will also be supercommutative, i.e., for homogeneous elements a and b ,

$$ab = (-1)^{|a||b|}ba,$$

where $|\cdot|$ is the \mathbb{Z}_2 -degree. Our algebras will furthermore be connected. Prime examples from this class of algebras are the polynomial ring $k[x_1, \dots, x_n]$, where $\deg x_i = (1, 0)$, and the exterior algebra $E(x_1, \dots, x_n)$, where $\deg x_i = (1, 1)$, and graded quotients of these.

The reason for considering this class is that Tate and Gulliksen's [Tat57, Gul68] construction of minimal resolutions of k which have a DGA structure applies. If W is a DGA, we will use the standard notation

$$\tilde{Z}(W) = \operatorname{Ker}(Z(W) \rightarrow k),$$

$$\tilde{H}(W) = \operatorname{Ker}(H(W) \rightarrow k),$$

$$IW = \operatorname{Ker}(W \rightarrow k).$$

For an \mathbb{N}^n -graded algebra A , and \mathbb{N}^n -graded A -module M , the Poincaré series of M is a formal power series defined by

$$P_A^M(y_1, \dots, y_n, z) = P_A^M(\mathbf{y}, z) = \sum_{i,j} \dim_k \operatorname{Ext}_A^{i,j}(M, k) \mathbf{y}^j z^i.$$

Note that the \mathbb{Z}_2 -grading is not taken into account in this definition.

DEFINITION 1. Let R be an $\mathbb{N} \times \mathbb{Z}_2$ -graded connected supercommutative k -algebra, and I be a homogeneous ideal and let $f: R \rightarrow R/I = \bar{R}$ be the natural projection. Then f is said to be a *Golod homomorphism* if the following condition is satisfied:

$$P_R^k(y, z) = \frac{P_R^k(y, z)}{1 - z^2 P_R^I(y, z)}.$$

Remark 1. The definition is from Levin [Lev76], who assumed R to be a commutative local ring. The situation in the supercommutative situation is a special case of the definition of Golod homomorphisms of DGA:s given by Avramov [Avr86]. Levin also gives the following sufficient condition for a projection $R \rightarrow R/I$ to be a Golod homomorphism: let X be a minimal DGA-resolution of k over R , then, according to [Lev76, Lemma 1.6], the existence of an R/I -submodule V of $X \otimes_R R_+/I$ such that $V^2 = 0$ and $\tilde{Z}(X \otimes_R R/I) \subset V + B(X \otimes_R R/I)$, implies that the projection is a Golod homomorphism. This condition will be used in the proof of Lemma 3.

DEFINITION 2. Let $I \subset S$ or $J \subset E$ be a homogeneous ideal. The quotient S/I or E/J is said to be a *Golod ring* if $S \rightarrow S/I$ or $E \rightarrow E/J$ is a Golod homomorphism.

3. THE MAIN RESULT

We will now state the main result of the paper.

THEOREM 1. *Let Δ be a finite simplicial complex; then $k[\Delta]$ is a Golod ring if and only if $k\{\Delta\}$ is a Golod ring.*

The proof depends on some lemmas that we will obtain before proving the theorem. The first of these lemmas is due to Aramova, Avramov, and Herzog; we will just reformulate it to suit our needs.

LEMMA 1. *The power series $P_E^{k[\Delta]}$ and $P_S^{k[\Delta]}$ are related by*

$$P_E^{k[\Delta]}(y_1, \dots, y_n, z) = P_S^{k[\Delta]} \left(\frac{y_1}{1 - y_1 z}, \dots, \frac{y_n}{1 - y_n z}, z \right).$$

Proof. For $i \in \mathbb{N}$ and $\mathbf{j} \in \mathbb{N}^n$, define $\beta_{i,\mathbf{j}}$ by $P_S^{k[\Delta]} = \sum_{i,\mathbf{j}} \beta_{i,\mathbf{j}} \mathbf{y}^{\mathbf{j}} z^i$. According to [AAH, Proposition 2.1],

$$P_E^{k[\Delta]} = \sum_{i,\mathbf{j}} \beta_{i,\mathbf{j}} \frac{\mathbf{y}^{\mathbf{j}}}{\prod_{\nu, j_\nu \neq 0} (1 - y_\nu z)} z^i.$$

The homology of $k[\Delta]$ is only nonzero in squarefree multidegrees, i.e., $\beta_{i,\mathbf{j}} = 0$ if any $j_\nu \notin \{0, 1\}$ (see Fröberg [Frö82]); this implies the desired formula. ■

The other main ingredient of the proof of Theorem 1 is to replace computations over $k\{\Delta\}$ with a series of computations over the rings $R_i = k[\Delta]/(x_1^2, \dots, x_i^2)$; we think of $k[\bar{\Delta}] = R_n$ as being the “Artinification” of $k[\Delta]$.

Let \mathcal{E} denote the category of multigraded $k[\bar{\Delta}]$ -modules, i.e., the objects are finitely generated \mathbb{N}^n -graded modules, and the morphisms are $k[\bar{\Delta}]$ -linear maps $f: M \rightarrow N$ such that $f(M_{\mathbf{j}}) \subset N_{\mathbf{j}}$, for $\mathbf{j} \in \mathbb{N}^n$. Similarly, let \mathcal{D} denote the category of multigraded left $k\{\Delta\}$ -modules.

Define a functor $F: \mathcal{E} \rightarrow \mathcal{D}$ in the following way. For a module M in \mathcal{E} define $F(M)$ by letting it have the same underlying k -space as M , and for a homogeneous element $m \in M$ of multidegree $(\alpha_1, \dots, \alpha_n)$ we define the $k\{\Delta\}$ -action $x_i \circ m$ on m by

$$x_i \circ m = (-1)^{\sum_{j < i} \alpha_j} x_i \bullet m,$$

where $x_i \bullet m$ is the $k[\bar{\Delta}]$ -action on m . It is now easy to see that this is indeed a $k\{\Delta\}$ -action. For a morphism ϕ we define $F(\phi) := \phi$.

Similarly, we may define another functor $G: \mathcal{D} \rightarrow \mathcal{E}$, by (for N in \mathcal{E}) letting $G(N)$ have the same k -structure as N , and define

$$x_i \bullet n = (-1)^{\sum_{j < i} \alpha_j} x_i \circ n$$

for $n \in N$ a homogeneous element of multidegree (a_1, \dots, a_n) . As before $G(\psi) = \psi$.

PROPOSITION 1. *The functors F and G defined above are additive, exact, mutually quasi-inverse equivalences of categories that commute with Exts.*

Proof. It is clear that F and G are additive, exact, and quasi-inverse. If \mathbf{P} is a multigraded $k[\bar{\Delta}]$ -free resolution of K , then $F(\mathbf{P})$ is a multigraded $k\{\Delta\}$ -free resolution of $F(K)$, and the complexes of multigraded vector

spaces $\text{Hom}_{k[\Delta]}(F(\mathbf{P}), F(L))$ and $\text{Hom}_{k[\bar{\Delta}]}(\mathbf{P}, L)$ are equal. This implies an equality of Exts. ■

As an immediate consequence of the proposition, we obtain the following lemma.

LEMMA 2. *The following equality holds*

$$\mathbf{P}_{k[\Delta]}^k(\mathbf{y}, z) = \mathbf{P}_{k[\bar{\Delta}]}^k(\mathbf{y}, z).$$

Remark 2. Since the element $x_{i+1} \in R_i$ is a nonzerodivisor on (x_{i+1}) we get that $\mathbf{P}_{R_i}^{(x_{i+1})} = y_{i+1} \mathbf{P}_{R_i}^{(x_{i+1})}$.

LEMMA 3. *The canonical projection $R_i \rightarrow R_{i+1}$ is a Golod homomorphism, and*

$$\frac{1}{\mathbf{P}_{R_{i+1}}^k} = \frac{1 - y_{i+1} z^2 \mathbf{P}_{R_i}^{(x_{i+1})}}{\mathbf{P}_{R_i}^k}.$$

Proof. For the first statement we will mimic the proof of [Lev76, Theorem 2.3], and use the condition mentioned in Remark 1. The second statement then follows from Remark 2. Let X be a minimal DGA-resolution of k over R , consider the natural epimorphism $\eta: X \rightarrow X \otimes_R R/(x_{i+1}^2)$, and pick $z \in \tilde{Z}(X \otimes_R R/(x_{i+1}^2))$. Choose z' such that $\eta(z') = z$. Then $d(z') \in x_{i+1}^2 X$, and we write $d(z') = x_{i+1}^2 u$. Apply d once more, and we get

$$0 = d^2(z') = x_{i+1} d(x_{i+1} u).$$

By Remark 2 we get that $d(x_{i+1} u) = 0$. Let $T_{i+1} \in X_1$ such that $d(T_{i+1}) = x_{i+1}$, then $d(z' - T_{i+1} x_{i+1} u) = 0$, so since X is acyclic, we have a v such that $z' = T_{i+1} u + d(v)$. Thus $z = \eta(z') = \eta(T_{i+1}) \eta(x_{i+1} u) + d\eta(v)$. Choose V by $V = \eta(T_{i+1})(X \otimes_R R/(x_{i+1}^2))$; since $T_{i+1}^2 = 0$, V satisfies the required condition, and the map $R_i \rightarrow R_{i+1}$ is Golod. ■

LEMMA 4. *We have the identity*

$$\mathbf{P}_{k[\bar{\Delta}]}^k(y_1, \dots, y_n, z) = \mathbf{P}_{k[\Delta]}^k\left(\frac{y_1}{1 - y_1 z}, \dots, \frac{y_n}{1 - y_n z}, z\right).$$

Proof. Set $A = R_i/(x_{i+1})$, so that $\mathbf{P}_A^k = 1 + z \mathbf{P}_{R_i}^{(x_{i+1})}$. Since there are ring homomorphisms $A \rightarrow R_i \rightarrow A$ whose composition is the identity, we

have $P_{R_i}^k = P_{R_i}^A P_A^k$ by Herzog [Her77]. Using Lemma 3 we then get

$$\frac{1}{P_{R_{i+1}}^k} = \frac{1 - y_{i+1}z^2 P_{R_i}^{(x_{i+1})}}{P_{R_i}^k} = \frac{1 + y_{i+1}z - y_{i+1}z P_{R_i}^A}{P_{R_i}^k} = \frac{1 + y_{i+1}z}{P_{R_i}^k} - \frac{y_{i+1}z}{P_A^k}.$$

Note that $R_{i+1} \simeq A[x_{i+1}]/x_{i+1}I_\Gamma$, where Γ is the simplicial complex on the vertex set $[n] \setminus \{i+1\}$ defined by $\sigma \in \Gamma$ if and only if $\sigma \cup \{i+1\} \in \Delta$. From the proof of Backelin [Bac82, Lemma 1] we know that

$$\frac{1}{P_{R_i}^k} = \frac{1}{P_A^k} - \frac{z(1 + zP_A^{I_\Gamma})}{P_A^k} \cdot \frac{y_{i+1}}{1 + y_{i+1}z}. \quad (2)$$

Using this formula to eliminate $P_{R_i}^k$ from the expression for $1/P_{R_{i+1}}^k$, we get

$$\frac{1}{P_{R_{i+1}}^k} = \frac{1}{P_A^k} - \frac{z(1 + zP_A^{I_\Gamma})}{P_A^k} \cdot y_{i+1}. \quad (3)$$

Notice that the power series P_A^k and $P_A^{I_\Gamma}$ do not involve the variable y_{i+1} . Thus, replacing $y_{i+1}/(1 - y_{i+1}z)$ transforms the right-hand side of (2) into that of (3). This establishes the equality

$$P_{R_{i+1}}^k(y_1, \dots, y_n, z) = P_{R_i}^k\left(y_1, \dots, y_i, \frac{y_{i+1}}{1 - y_{i+1}z}, y_{i+2}, \dots, y_n, z\right).$$

Applying it successively with $i = 0, \dots, n-1$ we get the assertion of the lemma. ■

Proof of the theorem. Assume that $k[\Delta]$ is Golod; this is by definition equivalent to the equality

$$P_{k[\Delta]}^k(\mathbf{y}, z) = \frac{P_S^k(\mathbf{y}, z)}{1 - z(P_S^{k[\Delta]}(\mathbf{y}, z) - 1)}.$$

According to Lemma 1, Lemma 2, and Lemma 4, this implies that

$$\begin{aligned} P_{k[\Delta]}^k(\mathbf{y}, z) &= P_{k[\Delta]}^k\left(\frac{y_1}{1 - y_1z}, \dots, \frac{y_n}{1 - y_nz}, z\right) \\ &= \frac{P_S^k(y_1/(1 - y_1z), \dots, y_n/(1 - y_nz), z)}{1 - z(P_S^{k[\Delta]}(y_1/(1 - y_1z), \dots, y_n/(1 - y_nz), z) - 1)} \\ &= \frac{P_E^k(\mathbf{y}, z)}{1 - z(P_E^{k[\Delta]}(\mathbf{y}, z) - 1)}, \end{aligned}$$

so $k\{\Delta\}$ is Golod as well. By using that $y \mapsto y/(1 - yz)$ is invertible, with inverse $y \mapsto y/(1 + yz)$, we see that the converse is also true. ■

REFERENCES

- [AAH] A. Aramova, L. L. Avramov, and J. Herzog, Resolutions of monomial ideals and cohomology over exterior algebras, *Trans. Amer. Math. Soc.* to appear.
- [AHH97] A. Aramova, J. Herzog, and T. Hibi, Gotzmann theorems for exterior algebras and combinatorics, *J. Algebra* **191** (1997), 174–211.
- [AHH98] A. Aramova, J. Herzog, and T. Hibi, Squarefree lexsegment ideals, *Math. Z.* **228**, No. 2 (1998), 353–378.
- [Avr86] L. L. Avramov, Golod homomorphisms, in “Algebra, Algebraic Topology and Their Interactions (Stockholm, 1983),” Lecture Notes in Mathematics, Vol. 1183, pp. 59–78, Springer-Verlag, Berlin, 1986.
- [Bac82] J. Backelin, Les anneaux locaux à relations monomiales ont des séries de Poincaré–Betti rationnelles, *C. R. Acad. Sci. Paris Sér. I Math.* **295**, No. 11 (1982), 607–610.
- [Frö82] R. Fröberg, A study of graded extremal rings and of monomial rings, *Math. Scand.* **51** (1982), 22–34.
- [Gul68] T. H. Gulliksen, A proof of the existence of minimal R-algebra resolutions, *Acta Math.* **120** (1968), 53–58.
- [Her77] J. Herzog, Algebra retracts and Poincaré-series, *Manuscripta Math.* **21**, No. 4 (1977), 307–314.
- [Lev76] G. Levin, “Lectures on Golod homomorphisms,” Mat. Inst., Stockholm University, Vol. 15, 1976.
- [Pee96] I. Peeva, 0-Borel fixed ideals, *J. Algebra* **184** (1996), 945–984.
- [Tat57] J. Tate, Homology of Noetherian rings and local rings, *Illinois J. Math.* **1** (1957), 14–27.